

Literatur

- [1] Hamza, I.A., Jurzik, J., Stang, A., Sure, K., Uberla K., Wilhelm M. (2009a): Detection of human viruses in rivers of a densely-populated area in Germany using a virus adsorption elution method optimized for PCR analyses. *Water Res.* 43:2657-2668.
- [2] Hamza, I.A., Jurzik, L., K. Uberla, K., Wilhelm M. (2011): Evaluation of pepper mild mottle virus, human picobirnavirus and Torque teno virus as indicators of fecal contamination in river water. *Water Res.* 45:1358-1368.
- [3] Hamza, I.A., Jurzik, L., Wilhelm, M., Uberla, K. (2009b): Detection and quantification of human bocavirus in river water. *J Gen Virol.* 90:2634-2637.
- [4] Jurzik, L., Hamza, I.A., Puchert, W., Uberla, K., Wilhelm, M. (2010): Chemical and microbiological parameters as possible indicators for human enteric viruses in surface water. *Int J Hyg Environ Health.* 213:210-216.
- [5] Selinka, H.C., Botzenhart K., Feuerpfeil, I., Puchert, W., Schmoll, O., Szewzyk R., Willmitzer, H. (2011): Detection of viruses in raw water as a basic tool for risk assessment. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz.* 54:496-504.
- [6] Timm, C., S. Luther, L. Jurzik, I.A. Hamza, and T. Kistemann. 2016. Applying QMRA and DALY to assess health risks from river bathing. *Int J Hyg Environ Health.*
- [7] Dieter, H., Chorus, I., Krüger, W., Mendel, B. (2017): *Trinkwasser aktuell.* Erich Schmidt Verlag.
- [8] RWW: Das Mülheimer Verfahren (2012). RWW Broschüre. <http://www.rww.de/Trinkwasser/wasseraufbereitung/muelheimer-verfahren/>
- [9] Hohlmann, W., Schöpel, M., Konrads, B., Losen H. (2005): Betriebsoptimierung eines Wasserwerkes durch nachhaltige Ertüchtigung der Langsandsandfiltration, *energie/wasser-praxis* 6/2006, 20-26.
- [10] Leifels, M., Jurzik, L., Wilhelm, M., Hamza, I.A. (2015): Use of ethidium monoazide and propidium monoazide to determine viral infectivity upon inactivation by heat, UV-exposure and chlorine. *Int J Hyg Environ Health.* 218:686-693.
- [11] Tondera, K., K. Klaer, S. Roder, I. Brueckner, M. Strathmann, T. Kistemann, C. Schreiber, and J. Pinnekamp. 2015. Developing an easy-to-apply model for identifying relevant pathogen pathways into surface waters used for recreational purposes. *Int J Hyg Environ Health.*
- [12] Malzer, H.J., T. Aus der Beek, S. Muller, and J. Gebhardt. 2016. Comparison of different model approaches for a hygiene early warning system at the lower Ruhr River, Germany. *Int J Hyg Environ Health.* 219:671-680.
- [13] DWD. 2016. Deutscher Wetterdienst, www.dwd.de.
- [14] Strathmann, M., M. Horstkott, C. Koch, U. Gayer, and J. Wingender. 2016. The River Ruhr - an urban river under particular interest for recreational use and as a raw water source for drinking water: The collaborative research project „Safe Ruhr“ - microbiological aspects. *Int J Hyg Environ Health.* 219:643-661.
- [15] Rodriguez-Lazaro, D., N. Cook, F.M. Ruggeri, J. Sellwood, A. Nasser, M.S. Nascimento, M. D'Agostino, R. Santos, J.C. Saiz, A. Rzezutka, A. Bosch, R. Girones, A. Carducci, M. Muscillo, K. Kovac, M. Diez-Valcarce, A. Vantarakis, C.H. von Bonsdorff, A.M. de Roda Husman, M. Hernandez, and W.H. van der Poel. 2012. Virus hazards from food, water and other contaminated environments. *FEMS Microbiol Rev.* 36:786-814.
- [16] Bae, S., and S. Wuertz. 2012. Survival of host-associated bacteroidales cells and their relationship with *Enterococcus* spp., *Campylobacter jejuni*, *Salmonella enterica* serovar Typhimurium, and adenovirus in freshwater microcosms as measured by propidium monoazide-quantitative PCR. *Appl Environ Microbiol.* 78:922-932.
- [17] Quijada, N.M., G. Fongaro, C.R. Barardi, M. Hernandez, and D. Rodriguez-Lazaro. 2016. Propidium Monoazide Integrated with qPCR Enables the Detection and Enumeration of Infectious Enteric RNA and DNA Viruses in Clam and Fermented Sausages. *Front Microbiol.* 7:2008.
- [18] Leifels, M., I.A. Hamza, M. Krieger, M. Wilhelm, M. Mackowiak, and L. Jurzik. 2016. From Lab to Lake - Evaluation of Current Molecular Methods for the Detection of Infectious Enteric Viruses in Complex Water Matrices in an Urban Area. *Plos One.* 11.
- [19] Prevost, B., M. Goulet, F.S. Lucas, M. Joyeux, L. Moulin, and S. Wurtzer. 2016. Viral persistence in surface and drinking water: Suitability of PCR pre-treatment with intercalating dyes. *Water Res.* 91:68-76.
- [20] Heim, A., Ebnet C., Harste, G., Pring-Akerblom, P. (2003): Rapid and quantitative detection of human adenovirus DNA by real-time PCR. *J Med Virol.* 70:228-239.
- [21] Logan, C., O'Leary, J.J., O'Sullivan, N. (2006): Real-time reverse transcription-PCR for detection of rotavirus and adenovirus as causative agents of acute viral gastroenteritis in children. *J Clin Microbiol.* 44:3189-3195.
- [22] Tajima, M., Kotani, Y., Kurosawa, T., Miyasaka, M. (2013): Pitfalls in mouse norovirus (MNV) detection in fecal samples using RT-PCR, and construction of new MNV-specific primers. *Exp Anim.* 62:127-135.
- [23] Price, J.A., Yamanashi, W., McGee, J.M. (1992): Bacteriophage phi X-174 as an aerobiological marker for surgical plume generated by the electromagnetic field focusing system. *The Journal of hospital infection.* 21:39-50.